Indiana adolescent reproductive health facts

Indiana was ranked 20 out of 51 (50 states + the District of Columbia) on 2011 final teen birth rates among females aged 15-19 (with 1 representing the highest rate and 51 representing the lowest rate).[RH1] On a similar scale – where 1 is the highest teen *pregnancy* rate and 51 is the lowest – Indiana was ranked 33 out of 51 (50 states + the District of Columbia) in pregnancies to females aged 15- 19 in 2008.[RH2]

Number of births to females under 20 years of age, 2011 [RH1]

Total[1]	Indiana	United States
Females under 20 years of age	8,010	333,746
Females aged	**Indiana**	**United States**
Under 15	93	3,974
15-17	2,132	95,538
18-19	5,785	234,234
15-19	7,917	329,772
Mother's race/ethnicity	**Indiana**	**United States**
Non-Hispanic white	5,566	130,198
Non-Hispanic black	1,531	79,936
American Indian or Alaska Native[2,3]	*	6,897
Asian or Pacific Islander[2,3]	42	5,773
Hispanic[4]	832	111,236

[1] Includes all births, including those with Hispanic origin not stated and not shown separately.
[2] Race and Hispanic origin are reported separately on birth certificates. Persons of Hispanic origin may be of any race. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards.
[3] Includes persons of Hispanic origin according to mother's reported race.
[4] Includes all persons of Hispanic origin of any race.

Teen birth rate (births per 1,000 females aged 15–19), 2011 [RH1]

Total	Indiana	United States
Females aged 15-19	34.8	31.3
Females aged	**Indiana**	**United States**
15-17	16.0	15.4
18-19	61.2	54.1

Percent change in the teen birth rate, 1991-2011 and 2010-2011 [RH1, RH3, RH4]

Total	Indiana	United States
Change in rate to females aged 15-19 (1991[◊] to 2011)	-42%	-49%
Change in rate to females aged 15-19 (2010 to 2011)	-7%	-8%

[◊]The U.S. teen birth rate peaked at 61.8 births per 1,000 teen females aged 15-19 in 1991.

Of all births to females under 20 years of age, percent by race/ethnicity, 2011 [RH1]

Mother's race /ethnicity[1]	Indiana	United States
Non-Hispanic white	69%	39%
Non-Hispanic black	19%	24%
American Indian or Alaska Native[2,3]	*	2%
Asian or Pacific Islander[2,3]	1%	2%
Hispanic[4]	10%	33%

[1]Includes all births, including those with Hispanic origin not stated and not shown separately.
[2]Race and Hispanic origin are reported separately on birth certificates. Persons of Hispanic origin may be of any race. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards.
[3]Includes persons of Hispanic origin according to mother's reported race.
[4]Includes all persons of Hispanic origin of any race.

Percent repeat births, of all births to females under 20 years of age, 2011 [RH1]

Total[1]	Indiana	United States
Females under 20 years of age	16%	18%
Mother's race/ethnicity	**Indiana**	**United States**
Non-Hispanic white	15%	14%
Non-Hispanic black	19%	20%
American Indian or Alaska Native[2,3]	*	19%
Asian or Pacific Islander[2,3]	*	17%
Hispanic[4]	23%	20%

[1] Includes all births, including those with Hispanic origin not stated and not shown separately.
[2] Race and Hispanic origin are reported separately on birth certificates. Persons of Hispanic origin may be of any race. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards.
[3] Includes persons of Hispanic origin according to mother's reported race.
[4] Includes all persons of Hispanic origin of any race.

Of all births to females under 20 years of age, percent outside of marriage (nonmarital), 2011 [RH1]

Total	Indiana	United States
Females under 20 years of age	90%	89%
Females aged	**Indiana**	**United States**
Under 15	99%	99%
15-17	97%	95%
18-19	87%	86%
15-19	90%	88%

Percent low birthweight[§] births, among females under 20 years of age, 2011 [RH1]

Total	Indiana	United States
Females under 20 years of age	10.2%	9.6%

[§] Low birth weight is defined as less than 2,500 grams (about 5.5 pounds or 5 pounds and 8 ounces).

Teen pregnancy rate (pregnancies per 1,000 females aged 15-19), 2008 [RH2]

Total	Indiana	United States
Females aged 15-19	59	68
Females aged	**Indiana**	**United States**
15-17	29	37
18-19	100	113

Percent change in the teen pregnancy rate, 1988-2008 [RH2]

Total	Indiana	United States
Change in rate to females aged 15-19 (1988♦ to 2008)	-34%	-39%

♦ The teen pregnancy rate was 89 pregnancies per 1,000 females aged 15-19 in Indiana in 1988, and 111 in the U.S. The U.S. teen pregnancy rate peaked in 1990 at 116.9 pregnancies per 1,000 females aged 15-19, but state data are not available for that year.

Teen abortion rate (abortions per 1,000 females aged 15-19), 2008 [RH2]

Total	Indiana	United States
Females aged 15-19	8	18
Females aged	**Indiana**	**United States**
15-17	5	10
18-19	13	29

Percent change in the teen abortion rate, 1988-2008 [RH2]

Total	Indiana	United States
Change in rate to females aged 15-19 (1988♦♦ to 2008)	-68%	-58%

♦♦ The U.S. teen abortion rate peaked at 43.5 abortions per 1,000 females aged 15-19 in 1988. The teen abortion rate for Indiana was 25 abortions per 1,000 females aged 15-19 in 1988.

Sexual behaviors among high school students (grades 9-12), 2011 [RH5]

Percent of high school students who:	Indiana	United States
Have ever had sexual intercourse		
Total	51%	47%
Male	51%	49%
Female	51%	46%
Had sexual intercourse for the first time before 13 years of age		
Total	5%	6%
Male	7%	9%
Female	4%	3%
Have had sexual intercourse with 4 or more persons		
Total	17%	15%
Male	18%	18%
Female	15%	13%
Drank alcohol or used drugs before last sexual intercourse		
Total	20%	22%
Male	20%	26%
Female	20%	18%

HIV/AIDS education among high school students (grades 9-12), 2011[RH5]

Percent of high school students who:	Indiana	United States
Were ever taught in school about AIDS or HIV infection		
Total	90%	84%
Male	89%	84%
Female	90%	84%

Birth control use among sexually active high school students (grades 9-12), 2011[RH5]

Percent of high school students who:	Indiana	United States
Used a condom during last sexual intercourse		
Total	58%	60%
Male	61%	67%
Female	55%	54%
Used birth control pills before last sexual intercourse		
Total	27%	18%
Male	25%	13%
Female	28%	23%
Used Depo-Provera, Nuva Ring, Implanon, or any IUD before last sexual intercourse[¤]		
Total	7%	5%
Male	4%	3%
Female	11%	7%
Used both a condom during last sexual intercourse and birth control pills or Depo-Provera, Nuva Ring, Implanon, or any IUD before last sexual intercourse[Ж]		
Total	18%	9%
Male	14%	7%
Female	21%	12%
Did not use any method to prevent pregnancy during last sexual intercourse		
Total	13%	13%
Male	13%	11%
Female	13%	15%

¤ Exact question wording: Used Depo-Provera (or any injectable birth control), Nuva Ring (or any birth control ring), Implanon (or any implant), or any IUD before last sexual intercourse.

Ж Exact question wording: Used both a condom during last sexual intercourse and Depo-Provera (or any injectable birth control), Nuva Ring (or any birth control ring), Implanon (or any implant), or any IUD before last sexual intercourse.

Relationship violence and forced sex among high school students (grades 9-12), 2011[RH5]

Percent of high school students who:	Indiana	United States
Have been hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend (during the 12 months before the survey)		
Total	11%	9%
Male	12%	9%
Female	11%	9%
Were ever physically forced to have sexual intercourse (when they did not want to)		
Total	10%	8%
Male	5%	4%
Female	15%	12%

Data notes and data sources

All percentages (other than percent low birthweight births), have been rounded to the nearest whole number.

* Figure does not meet standards of reliability or precision; based on fewer than 20 births in the numerator.

N/A: data do not exist on this measure for this state.

[1] Includes all births, including those with Hispanic origin not stated and not shown separately.

[2] Race and Hispanic origin are reported separately on birth certificates. Persons of Hispanic origin may be of any race.
Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards.

[3] Includes persons of Hispanic origin according to mother's reported race.

[4] Includes all persons of Hispanic origin of any race.

[RH1] **2011 Birth data are from:** Centers for Disease Control and Prevention (2013). National Vital Statistics System. Hyattsville, MD: National Center for Health Statistics.

[RH2] **Pregnancy and abortion rate data are from:** Kost ,K. and Henshaw, S. (2013) *U.S. Teenage Pregnancies, Births and Abortions, 2008: State Trends by Age, Race and Ethnicity*, New York: Guttmacher Institute. Retrieved August 7, 2013, from http://www.guttmacher.org/pubs/USTPtrendsState08.pdf

[RH3] **1991 Birth rate data are from:** Martin, J. A., Hamilton, B. E., Ventura, S. J., Osterman, M. J. K., Kirmeyer, S., Mathews, T. J., Wilson, E. (2011). *Births: Final data for 2009. Table B.* National vital statistics reports 60(1). Hyattsville, MD: National Center for Health Statistics. Retrieved August 7, 2013, from http://www.cdc.gov/nchs/data/nvsr/nvsr60/nvsr60_01.pdf

[RH4] **2010 Birth rate data are from:** Martin, J. A., Hamilton, B. E., Ventura, S. J., Osterman, M. J. K., Wilson, E.C., and Mathews, T. J. (2012). *Births: Final data for 2010*. National vital statistics reports 61(1). Hyattsville, MD: National Center for Health Statistics. Retrieved August 7, 2013, from http://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_01.pdf

[RH5] **High school data are from:** Centers for Disease Control and Prevention. (2012). *1991-2011 High School Youth Risk Behavior Survey Data.* Retrieved August 7, 2013, from http://apps.nccd.cdc.gov/youthonline